Courant Institute of
Mathematical Sciences

Abel's Equation and the Cauchy
Integral Equation of the Second Kind

A. S. Peters

Prepared under Contract Nonr-285(55)
with the Office of Naval Research NR 062-160

Distribution of this document is unlimited.

New York University

NEW YORK UNIVERSITY
COURANT INSTITUTE - LIBRARY
251 Mercer St., New York, N.Y. 10012

New York University
Courant Institute of Mathematical Sciences

ABEL'S EQUATION AND THE CAUCHY INTEGRAL
EQUATION OF THE SECOND KIND

A. S. Peters

This report represents results obtained at the Courant
Institute of Mathematical Sciences, New York University,
with the Office of Naval Research, Contract Nonr-285(55).
Reproduction in whole or in part is permitted for any
purpose of the United States Government.

Distribution of this document is unlimited.

NEW YORK UNIVERSITY
COURANT INSTITUTE - LIBRARY

1. Introduction

Let $\phi(z) \equiv \phi(x+iy)$ be a complex function which is integrable along a simple smooth path L in the complex z -plane. Let $\zeta = \xi+i\eta$ be a point on L which is not one of the end points of L and let ζ be imbedded in a subarc L_ϵ of L such that the end points of L_ϵ are equidistant from ζ . In this paper the symbol

$$\int_L \frac{\phi(z)dz}{z-\zeta}$$

means the Cauchy principal value, that is,

$$\int_L \frac{\phi(z)dz}{z-\zeta} = \lim_{\epsilon \rightarrow 0} \int_{L-L_\epsilon} \frac{\phi(z)dz}{z-\zeta}$$

provided the limit exists.

In a recent note [1] the author showed that the solution of the integral equation

$$(1.1) \quad \int_0^1 \frac{\phi(x)dx}{x-\xi} = f(\xi) \quad , \quad 0 < \xi < 1$$

can be reduced to the solution of

$$(1.2) \quad \int_0^x \frac{1}{\sqrt{x-t}} \int_t^1 \frac{\sqrt{\xi} \phi(\xi) d\xi dt}{\sqrt{\xi-t}} = \int_0^x \sqrt{\xi} f(\xi) d\xi + 2c \sqrt{x}$$

and therefore to the solution of the pair of Volterra equations

$$(1.3) \quad \int_0^x \frac{\psi(t)dt}{\sqrt{x-t}} = \int_0^x \sqrt{\xi} f(\xi)d\xi + 2c \sqrt{x}$$

$$(1.4) \quad \int_t^1 \frac{\sqrt{\xi} \phi(\xi)d\xi}{\sqrt{\xi-t}} = \psi(t) .$$

Equation (1.3) is Abel's equation and equation (1.4) can be reduced to the form (1.3) by using the transformation $\xi = 1 - u$; $\sigma = 1 - v$. The solution of (1.3) followed by the solution of (1.4) gives

$$(1.5) \quad \sqrt{\xi} \phi(\xi) = \frac{c}{\pi\sqrt{1-\xi}} - \frac{1}{\pi^2} \frac{d}{d\xi} \int_{\xi}^1 \frac{1}{\sqrt{t-\xi}} \int_0^t \frac{\sqrt{x} f(x)dxdt}{\sqrt{t-x}} .$$

The analysis in [1] also implies that the solution of (1.1) can be based on a knowledge of the integral

$$(1.6) \quad - \int_0^t \frac{1}{\sqrt{\xi(t-x)}} \cdot \frac{d\xi}{\xi-x} = \begin{cases} 0 & 0 < x < t \\ \frac{\pi}{\sqrt{x(x-t)}} & t < x \end{cases}$$

and the solution of just one Abel equation in the following way.

In order to allow $\phi(x)$ to possess a possible singularity at $x = 0$ let (1.1) be written in the form

$$(1.7) \quad \int_0^1 \frac{x\phi(x)dx}{x-\xi} = \xi f(\xi) + c$$

where

$$c = \int_0^1 \phi(x) dx .$$

The multiplication of both sides of (1.7) by $1/\sqrt{\xi(t-\xi)}$ followed by an integration gives

$$\int_0^t \frac{1}{\sqrt{\xi(t-\xi)}} \int_0^1 \frac{x\phi(x)dx d\xi}{x-\xi} = \int_0^t \frac{\sqrt{\xi} f(\xi)d\xi}{\sqrt{t-\xi}} + c \int_0^t \frac{d\xi}{\sqrt{\xi(t-\xi)}}$$

or

$$(1.8) \quad - \int_0^1 x\phi(x) \int_0^t \frac{1}{(\xi-x)\sqrt{\xi(t-\xi)}} d\xi dx = \int_0^t \frac{\sqrt{x} f(x)dx}{\sqrt{t-x}} + \pi c .$$

From (1.6) the equation (1.8) is the same as

$$\int_t^1 \frac{\sqrt{x} \phi(x)dx}{\sqrt{x-t}} = c + \frac{1}{\pi} \int_0^t \frac{\sqrt{x} f(x)dx}{\sqrt{t-x}} .$$

The solution of this integral equation is readily found to be

$$\sqrt{x} \phi(x) = \frac{c}{\pi\sqrt{1-x}} - \frac{1}{\pi^2} \frac{d}{dx} \int_x^1 \frac{1}{\sqrt{t-x}} \int_0^t \frac{\sqrt{x} f(x)dx dt}{\sqrt{t-x}}$$

which agrees with (1.5).

The method presented in the last paragraph depends on (1.6) which shows that

$$\psi(\xi) = \frac{1}{\sqrt{\xi(t-\xi)}}$$

is a solution of the homogeneous equation

$$\int_0^t \frac{\psi(\xi) d\xi}{\xi - x} = 0, \quad 0 < x < \xi.$$

More generally, it may be possible to base the solution of the Fredholm equation

$$(1.9) \quad \int_a^b K(\xi, x) \phi(x) dx = \lambda \phi(\xi) + f(\xi), \quad a < \xi < b$$

on the use of an explicit non-trivial solution $\psi(x, t)$ of

$$(1.10) \quad \int_a^t K(x, \xi) \psi(x, t) dx = \lambda \psi(\xi, t), \quad a < \xi < t.$$

If (1.9) is multiplied by $\psi(\xi, t)$ and then integrated it becomes

$$\int_a^t \psi(\xi, t) \int_a^b K(\xi, x) \phi(x) dx d\xi = \lambda \int_0^t \psi(\xi, t) \phi(\xi) d\xi + \int_0^t \psi(\xi, t) f(\xi) d\xi$$

or, if the order of integration can be changed,

$$\int_a^b \phi(x) \int_a^t K(\xi, x) \psi(\xi, t) d\xi dx = \lambda \int_0^t \psi(\xi, t) \phi(\xi) d\xi + \int_0^t \psi(\xi, t) f(\xi) d\xi$$

which is the same as

$$\int_a^t \phi(x) \int_a^t K(\xi, x) \psi(\xi, t) d\xi dx + \int_t^b \phi(x) \int_a^t K(\xi, x) \psi(\xi, t) d\xi dx$$

$$= \lambda \int_a^t \psi(\xi, t) \phi(\xi) d\xi + \int_a^t \psi(\xi, t) f(\xi) d\xi .$$

From (1.10) this is equivalent to

$$(1.11) \quad \int_t^b \phi(x) \int_a^t K(\xi, x) \psi(\xi, t) d\xi dx = \int_a^t \psi(\xi, t) f(\xi) d\xi .$$

If $K_1(t, x)$ denotes the value of the integral

$$\int_a^t K(\xi, x) \psi(\xi, t) d\xi$$

for $x > t$ the equation (1.11) becomes

$$(1.12) \quad \int_t^b K_1(t, x) \phi(x) dx = \int_a^t \psi(\xi, t) f(\xi) d\xi .$$

Therefore if a non-trivial solution of (1.10) can be found for $a < t < b$ then it may be effective to reduce the solution of the Fredholm equation (1.9) in the above way to the solution of the Volterra equation (1.12).

When the kernel $K(\xi, x)$ is a Cauchy kernel, that is, when $K(\xi, x) = 1/(x - \xi)$ the method just presented can be used to extend the results of [1]. Section 2, below, shows that the above method can be used to reduce the solution of

$$(1.13) \quad \int_0^1 \frac{\phi(x)dx}{x-\xi} = \lambda\phi(\xi) + f(\xi)$$

to the solution of Abel's integral equation. Section 3 demonstrates that the same ideas can be applied to obtain the solution of

$$(1.14) \quad \int_L \frac{\phi(z)dz}{z-\xi} = \lambda\phi(\xi) + f(\xi) .$$

In other words, the evaluation of a certain definite integral can be used to reduce the solution of (1.14) to the solution of a generalized Abel's equation. This leads to a new formula for the solution of (1.14).

2. Abel's Equation and a Cauchy Integral Equation of the Second Kind

The integral (1.6), namely

$$- \int_0^t \frac{d\xi}{(\xi-x)\sqrt{\xi(t-\xi)}}$$

is a special case of

$$- \int_0^t \left(\frac{t-\xi}{\xi}\right)^\gamma \frac{d\xi}{(t-\xi)(\xi-x)}$$

where $0 < \gamma < 1$. The substitution $t - \xi = \sigma\xi$ changes the last integral into

$$\int_0^{\infty} \frac{\sigma^{\gamma-1} dx}{x\sigma - (t-x)} d\sigma$$

which can be evaluated in a variety of ways. In one way or another it is not difficult to verify that

$$(2.1) \quad - \int_0^t \left(\frac{t-\xi}{\xi}\right)^{\gamma} \frac{d\xi}{(t-\xi)(\xi-x)}$$

$$= \begin{cases} \frac{1}{x} \left(\frac{t-x}{x}\right)^{\gamma-1} \int_0^{\infty} \frac{\xi^{\gamma-1} d\xi}{\xi-1} = - \frac{(t-x)^{\gamma-1}}{x^{\gamma}} \pi \cot \gamma\pi, & 0 < x < t \\ \frac{1}{x} \left(\frac{t-x}{x}\right)^{\gamma-1} \int_0^{\infty} \frac{\xi^{\gamma-1} d\xi}{\xi+1} = \frac{(x-t)^{\gamma-1}}{x^{\gamma}} \cdot \frac{\pi}{\sin \gamma\pi}, & t < x. \end{cases}$$

This shows that if λ is a given real value and γ is chosen so that

$$(2.2) \quad -\pi \cot \gamma\pi = \lambda, \quad 0 < \gamma < 1$$

then

$$(2.3) \quad \psi(\xi, t) = \frac{(t-\xi)^{\gamma-1}}{\xi^{\gamma}}$$

is a solution of

$$(2.4) \quad - \int_0^t \frac{\psi(\xi, t) d\xi}{\xi-x} = \lambda \psi(x, t), \quad 0 < x < t.$$

Once this is recognized one can surmise that the solution of (2.4) can be used in various ways to deduce the solution of the nonhomogeneous equation

$$(2.5) \quad \int_0^1 \frac{\phi(x)dx}{x-\xi} = \lambda\phi(\xi) + f(\xi)$$

in which all quantities are supposed real. For example, (2.3) and (2.4) could be used in connection with the Hardy-Poincaré-Bertrand formula [2] to find the solution of (2.5) but the author believes that the simplest procedure is to use the method outlined in the introduction. In accordance with this method the solution (2.3) can be used to reduce (2.5) to a simple Volterra equation.

Equation (2.5) can be expressed in the form

$$(2.6) \quad \int_0^1 \frac{x\phi(x)dx}{x-\xi} = \lambda\xi\phi(\xi) + \xi f(\xi) + c$$

where

$$c = \int_0^1 \phi(x)dx .$$

Then, multiplication of (2.6) by $(t-\xi)^{\gamma-1}/\xi^\gamma$ and integration gives

$$(2.7) \quad \int_0^t \frac{(t-\xi)^{\gamma-1}}{\xi^\gamma} \int_0^1 \frac{x\phi(x)dx d\xi}{x-\xi} = \int_0^t \frac{(t-\xi)^{\gamma-1}}{\xi^\gamma} [\lambda\xi\phi(\xi) + \xi f(\xi) + c] d\xi .$$

Under the assumption that the order of integration can be changed,
(2.7) is

$$\begin{aligned}
 (2.8) \quad & - \int_0^t x \phi(x) \int_0^t \frac{(t-\xi)^{\gamma-1}}{\xi^\gamma} \cdot \frac{d\xi dx}{\xi-x} - \int_t^1 x \phi(x) \int_0^t \frac{(t-\xi)^{\gamma-1}}{\xi^\gamma} \cdot \frac{d\xi dx}{\xi-x} \\
 & = \int_0^t \frac{(t-\xi)^{\gamma-1}}{\xi^\gamma} [\lambda \xi \phi(\xi) + \xi f(\xi) + c] d\xi .
 \end{aligned}$$

With γ determined by (2.2), the evaluation (2.1) can be used to reduce (2.8) to

$$\begin{aligned}
 (2.9) \quad & \frac{\pi}{\sin \gamma \pi} \int_t^1 (x-t)^{\gamma-1} x^{1-\gamma} \phi(x) dx \\
 & = \int_0^t (t-\xi)^{\gamma-1} \xi^{1-\gamma} f(\xi) d\xi + c \int_0^t \frac{(t-\xi)^{\gamma-1} d\xi}{\xi^\gamma} .
 \end{aligned}$$

Thus, since

$$\int_0^t \frac{(t-\xi)^{\gamma-1} d\xi}{\xi^\gamma} = \frac{\pi}{\sin \gamma \pi} ,$$

the solution of (2.5) reduces to the solution of

$$(2.10) \quad \int_t^1 (x-t)^{\gamma-1} x^{1-\gamma} \phi(x) dx = \frac{\sin \gamma \pi}{\pi} \int_0^t (t-\xi)^{\gamma-1} \xi^{1-\gamma} f(\xi) d\xi + c .$$

Furthermore, it is not difficult to prove that the process can be reversed so that any solution of (2.10) is also a solution of (2.5).

The solution of Abel's integral equation

$$(2.11) \quad \int_0^t \frac{\psi_1(x)dx}{(t-x)^\kappa} = F(t) \quad , \quad 0 < \kappa < 1$$

is

$$(2.12) \quad \psi_1(x) = \frac{\sin \kappa \pi}{\pi} \frac{d}{dx} \int_0^x \frac{F(t)dt}{(x-t)^{1-\kappa}} \quad .$$

The equation

$$(2.13) \quad \int_t^1 \frac{\psi_2(x)dx}{(x-t)^\kappa} = F(t) \quad , \quad 0 < \kappa < 1$$

can be reduced to (2.11). The substitutions $x = 1 - u$; $t = 1 - v$ change (2.13) into

$$\int_0^v \frac{\psi_2(1-u)du}{(v-u)^\kappa} = F(1-v)$$

and then by using (2.12) the solution of (2.13) is

$$(2.14) \quad \psi_2(x) = - \frac{\sin \kappa \pi}{\pi} \frac{d}{dx} \int_x^1 \frac{F(t)dt}{(t-x)^{1-\kappa}} \quad .$$

The application of (2.14) to (2.10) leads to

$$(2.15) \quad x^{1-\gamma} \phi(x) = - \frac{\sin^2 \gamma \pi}{\pi^2} \frac{d}{dx} \int_x^1 \frac{1}{(t-x)^\gamma} \int_0^t (t-\xi)^{\gamma-1} \xi^{1-\gamma} f(\xi) d\xi dt \\ + \frac{c \sin \gamma \pi}{\pi(1-x)^\gamma} \quad .$$

Since

$$-\pi \cot \gamma\pi = \lambda$$

and consequently

$$\frac{\sin \gamma\pi}{\pi} = \frac{1}{\sqrt{\lambda^2 + \pi^2}} ;$$

the result (2.15) can also be expressed as

$$(2.16) \quad x^{1-\gamma}\phi(x) = - \frac{1}{\lambda^2 + \pi^2} \frac{d}{dx} \int_x^1 \frac{1}{(t-x)^\gamma} \int_0^t (t-\xi)^{\gamma-1} \xi^{1-\gamma} f(\xi) d\xi dt \\ + \frac{c}{(1-x)^\gamma \sqrt{\lambda^2 + \pi^2}} .$$

The formula (2.16) is not the formula which is usually given for the solution of (2.5). However, the derivative in (2.16) can be transformed as follows:

$$\frac{d}{dx} \int_x^1 \frac{1}{(t-x)^\gamma} \int_0^t (t-\xi)^{\gamma-1} \xi^{1-\gamma} f(\xi) d\xi dt \\ = \frac{d}{dx} \left\{ \int_0^x \xi^{1-\gamma} f(\xi) \int_x^1 \frac{(t-\xi)^{\gamma-1} dt d\xi}{(t-x)^\gamma} + \int_x^1 \xi^{1-\gamma} f(\xi) \int_\xi^1 \frac{(t-\xi)^{\gamma-1} dt d\xi}{(t-x)^\gamma} \right\} \\ = \frac{d}{dx} \left\{ \int_0^x \xi^{1-\gamma} f(\xi) \int_\infty^{\frac{1-\xi}{1-x}} \frac{\sigma^{\gamma-1} d\sigma d\xi}{1-\sigma} + \int_x^1 \xi^{1-\gamma} f(\xi) \int_0^{\frac{1-\xi}{1-x}} \frac{\sigma^{\gamma-1} d\sigma d\xi}{1-\sigma} \right\}$$

$$\begin{aligned}
&= x^{1-\gamma} f(x) \int_{\infty}^1 \frac{\sigma^{\gamma-1} d\sigma}{1-\sigma} - x^{1-\gamma} f(x) \int_0^1 \frac{\sigma^{\gamma-1} d\sigma}{1-\sigma} \\
&\quad + \int_0^1 \xi^{1-\gamma} f(\xi) \left(\frac{1-\xi}{1-x} \right)^{\gamma-1} \cdot \frac{1}{1 - \left(\frac{1-\xi}{1-x} \right)} \cdot \frac{(1-\xi) d\xi}{(1-x)^2} \\
&= x^{1-\gamma} f(x) \int_0^{\infty} \frac{\sigma^{\gamma-1} d\sigma}{\sigma-1} + \frac{1}{(1-x)^{\gamma}} \int_0^1 \frac{(1-\xi)^{\gamma} \xi^{1-\gamma} f(\xi) d\xi}{\xi-x} \\
&= -x^{1-\gamma} f(x) \pi \cot \gamma\pi + \frac{1}{(1-x)^{\gamma}} \int_0^1 \frac{(1-\xi)^{\gamma} \xi^{1-\gamma} f(\xi) d\xi}{\xi-x} \\
&= \lambda x^{1-\gamma} f(x) + \frac{1}{(1-x)^{\gamma}} \int_0^1 \frac{(1-\xi)^{\gamma} \xi^{1-\gamma} f(\xi) d\xi}{\xi-x} .
\end{aligned}$$

Therefore by using this transformation, which we will denote by T, equation (2.16) can be expressed as

$$\begin{aligned}
(2.17) \quad \phi(x) = & - \frac{\lambda f(x)}{\lambda^2 + \pi^2} - \frac{1}{(\lambda^2 + \pi^2)} \cdot \frac{1}{x^{1-\gamma} (1-x)^{\gamma}} \int_0^1 \frac{(1-\xi)^{\gamma} \xi^{1-\gamma} f(\xi) d\xi}{\xi-x} \\
& + \frac{c}{x^{1-\gamma} (1-x)^{\gamma} \sqrt{\lambda^2 + \pi^2}}
\end{aligned}$$

where

$$-\pi \cot \gamma\pi = \lambda$$

The formula (2.17) appears in Muskhelishvili [2] or Mikhlin [3], for instance, and it is regarded as the standard formula for the solution of

$$(2.18) \quad \int_0^1 \frac{\phi(x)dx}{x-\xi} = \lambda\phi(\xi) + f(\xi) \quad .$$

Incidentally, the solution (2.15) can be checked by using successive Abel equations to solve for $f(\xi)$ in (2.15) and then applying a transformation similar to the transformation T noted above.

3. The Cauchy Integral Equation of the Second Kind with a Path of Integration in the Complex Plane

If L is a simple smooth path from α to β in the complex z -plane and if ξ is on L' , which denotes L minus its end points, the equation

$$(3.1) \quad \int_L \frac{\phi(z)dz}{z-\xi} = \lambda\phi(\xi) + f(\xi)$$

is a generalization of (2.18). As is well known (see [2], [3], [4], [5]) the solution of (3.1) can be reduced to the solution of a Hilbert-Riemann boundary value problem. The reduction, as well as the solution of the subsidiary problem, is based on the use of an ingenious idea due to Carleman [6], namely the introduction of the function

$$F(w) = \int_L \frac{\phi(z)dz}{z-w}$$

where w is unrestricted; and the use of the Plemelj formulas. This method for finding the solution of (3.1) is very elegantly effective and it is perhaps indispensable for finding the integral representation for the solution of a Cauchy integral equation of the third kind. For these reasons and others, the method is likely to remain in its position of preeminence. However, apart from an analysis of the properties of $\phi(z)$ which will insure the existence of the Cauchy principal value for all or almost all values of ζ , it could be claimed that the ideas mentioned above require the development of a function theoretic apparatus which is unnecessarily advanced and powerful for the solution of the real equation (2.5) if one grants a knowledge of the integral (2.1). Compared with such a claim, a similar claim with respect to equation (3.1), which involves complex quantities, is less defensible. Nevertheless, the purpose of this section is to show that a solution of (3.1) analogous to that of (2.5) in Section 2 can be synthesized if one is willing to start with the value of a definite integral which is a generalization of (2.1). The generalized integral is

$$(3.2) \quad - \int_{L_{\alpha\omega}} \left(\frac{\omega-\zeta}{\zeta-\alpha} \right)^{\gamma} \frac{d\zeta}{(\omega-\zeta)(\zeta-z)} \quad , \quad 0 < \operatorname{Re}(\gamma) < 1$$

where $L_{\alpha\omega}$ is the path along L from α to an arbitrary point ω on L . With the evaluation of this integral as a basis, the solution of (3.1) can be reduced in an elementary way to the solution of a generalization of Abel's equation.

The substitution $\omega - \xi = \sigma(\xi - \alpha)$ changes (3.2) into the simpler form

$$(3.3) \quad \int_{\Gamma_{0\infty}} \frac{\sigma^{\gamma-1} d\sigma}{\sigma(z-\alpha) - (\omega-z)}$$

where $\Gamma_{0\infty}$ denotes a simple smooth path from the origin to infinity in the complex σ -plane. If z is on $L_{\alpha\omega}$, $(\omega-z)/(z-\alpha)$ is on $\Gamma_{0\infty}$; and if z is on $L-L_{\alpha\omega}$, $(\omega-z)/(z-\alpha)$ is not on $\Gamma_{0\infty}$. It is easy to evaluate (3.3) by using the theory of residues and this gives

$$(3.4) \quad - \int_{L_{\alpha\omega}} \left(\frac{\omega-\xi}{\xi-\alpha} \right)^{\gamma} \frac{d\xi}{(\omega-\xi)(\xi-z)}$$

$$= \begin{cases} - \left(\frac{\omega-z}{z-\alpha} \right)^{\gamma-1} \frac{\pi \cot \gamma\pi}{(z-\alpha)} & z \text{ on } L'_{\alpha\omega} \\ \left(\frac{z-\omega}{z-\alpha} \right)^{\gamma-1} \frac{\pi}{(z-\alpha) \sin \gamma\pi} & z \text{ on } L-L_{\alpha\omega} \end{cases} .$$

The evaluation (3.4) shows that if $\lambda (\neq \pm\pi i)$ is given and γ is chosen so that

$$(3.5) \quad -\pi \cot \gamma\pi = \lambda , \quad 0 < \operatorname{Re}(\gamma) < 1 ,$$

then

$$(3.6) \quad \psi(\xi, \omega) = \frac{(\omega-\xi)^{\gamma-1}}{(\xi-\alpha)^{\gamma}}$$

is a solution of the homogeneous equation

$$(3.7) \quad - \int_{L_{\alpha\omega}} \frac{\psi(\xi, \omega) d\xi}{\xi - z} = \lambda \psi(z, \omega)$$

where z is a point on $L'_{\alpha\omega}$.

Equation (3.1) can be written as

$$(3.8) \quad \int_L \frac{(z-\alpha)\phi(z)dz}{z-\xi} = \lambda(\xi-\alpha)\phi(\xi) + (\xi-\alpha)f(\xi) + c$$

where

$$c = \int_L \phi(z)dz .$$

Multiplication of (3.8) by $(\omega-\xi)^{\gamma-1}/(\xi-\alpha)^\gamma$ followed by an integration yields

$$(3.9) \quad \int_{L_{\alpha\omega}} \frac{(\omega-\xi)^{\gamma-1}}{(\xi-\alpha)^\gamma} \int_L \frac{(z-\alpha)\phi(z)dzd\xi}{z-\xi} \\ = \int_{L_{\alpha\omega}} \frac{(\omega-\xi)^{\gamma-1}}{(\xi-\alpha)^\gamma} [\lambda(\xi-\alpha)\phi(\xi) + (\xi-\alpha)f(\xi) + c]d\xi .$$

The assumption that the order of integration can be changed in (3.9) leads to

$$\begin{aligned}
(3.10) \quad & - \int_{L_{\alpha\omega}} (z-\alpha)\phi(z) \int_{L_{\alpha\omega}} \frac{(\omega-\xi)^{\gamma-1}}{(\xi-\alpha)^\gamma} \frac{d\xi dz}{\xi-z} \\
& - \int_{L_{\omega\beta}} (z-\alpha)\phi(z) \int_{L_{\alpha\omega}} \frac{(\omega-\xi)^{\gamma-1}}{(\xi-\alpha)^\gamma} \frac{d\xi dz}{\xi-z} \\
& = \int_{L_{\alpha\omega}} \frac{(\omega-\xi)^{\gamma-1}}{(\xi-\alpha)^\gamma} [\lambda(\xi-\alpha)\phi(\xi) + (\xi-\alpha)f(\xi) + c] d\xi
\end{aligned}$$

where $L_{\omega\beta}$ is the path along L from ω to β . If γ is chosen so that (3.5) holds then (3.4) shows that (3.10) is the same as

$$\begin{aligned}
(3.11) \quad & \frac{\pi}{\sin \gamma\pi} \int_{L_{\omega\beta}} \frac{(z-\alpha)^{1-\gamma}}{(z-\omega)^{1-\gamma}} \phi(z) dz \\
& = \int_{L_{\alpha\omega}} \frac{(\xi-\alpha)^{1-\gamma} f(\xi) d\xi}{(\omega-\xi)^{1-\gamma}} + c \int_{L_{\alpha\omega}} \frac{(\omega-\xi)^{\gamma-1} d\xi}{(\xi-\alpha)^\gamma} .
\end{aligned}$$

Since

$$(3.12) \quad \int_{L_{\alpha\omega}} \frac{(\omega-\xi)^{\gamma-1} d\xi}{(\xi-\alpha)^\gamma} = \frac{\pi}{\sin \gamma\pi}$$

equation (3.11) reduces to

$$(3.13) \quad \int_{L_{\omega\beta}} \frac{(z-\alpha)^{1-\gamma} \phi(z) dz}{(z-\omega)^{1-\gamma}} = \frac{\sin \gamma\pi}{\pi} \int_{L_{\alpha\omega}} \frac{(\xi-\alpha)^{1-\gamma} f(\xi) d\xi}{(\omega-\xi)^{1-\gamma}} + c$$

which may be classified as a generalized Abel equation. The equation can be solved by using (2.14). However, for completeness, a derivation of the solution follows.

Equation (3.13) can be solved by multiplying each side by $1/(\omega-\tau)^\gamma$ where τ is on $L_{\alpha\beta}$; and integrating along L from τ to β . This gives

$$\begin{aligned} \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\omega\beta}} \frac{(z-\alpha)^{1-\gamma} \phi(z) dz d\omega}{(z-\omega)^{1-\gamma}} \\ = \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\tau\beta}} \frac{(z-\alpha)^{1-\gamma} \mu(z-\omega) \phi(z) dz d\omega}{(z-\omega)^{1-\gamma}} \end{aligned}$$

where

$$\mu(z-\omega) = \begin{cases} 1 & \text{if } z \text{ follows } \omega \text{ along } L \\ 0 & \text{if } z \text{ precedes } \omega \text{ along } L. \end{cases}$$

Hence it can be seen that

$$\begin{aligned} \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\omega\beta}} \frac{(z-\alpha)^{1-\gamma} \phi(z) dz d\omega}{(z-\omega)^{1-\gamma}} \\ = \int_{L_{\tau\beta}} (z-\alpha)^{1-\gamma} \phi(z) \int_{L_{\tau\beta}} \frac{\mu(z-\omega) d\omega dz}{(\omega-\tau)^\gamma (z-\omega)^{1-\gamma}} \\ = \int_{L_{\tau\beta}} (z-\alpha)^{1-\gamma} \phi(z) \int_{L_{\tau z}} \frac{d\omega dz}{(\omega-\tau)^\gamma (z-\omega)^{1-\gamma}} \end{aligned}$$

or by using (3.12)

$$\int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\omega\beta}} \frac{(z-\alpha)^{1-\gamma} \phi(z) dz d\omega}{(z-\omega)^{1-\gamma}} = \frac{\pi}{\sin \gamma\pi} \int_{L_{\tau\beta}} (z-\alpha)^{1-\gamma} \phi(z) dz .$$

From this, the solution of (3.13) is given by

$$\begin{aligned} (3.14) \quad & \frac{\pi}{\sin \gamma\pi} \int_{L_{\tau\beta}} (z-\alpha)^{1-\gamma} \phi(z) dz \\ &= \frac{\sin \gamma\pi}{\pi} \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\alpha\omega}} \frac{(\xi-\alpha)^{1-\gamma} f(\xi) d\xi d\omega}{(\omega-\xi)^{1-\gamma}} + c \int_{L_{\tau\beta}} \frac{d\omega}{(\omega-\tau)^\gamma} . \end{aligned}$$

The derivative of (3.14) is

$$\begin{aligned} (3.15) \quad & (\tau-\alpha)^{1-\gamma} \phi(\tau) \\ &= - \frac{\sin^2 \gamma\pi}{\pi^2} \frac{d}{d\tau} \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\alpha\omega}} \frac{(\xi-\alpha)^{1-\gamma} f(\xi) d\xi d\omega}{(\omega-\xi)^{1-\gamma}} + \frac{c \sin \gamma\pi}{\pi(\beta-\tau)^\gamma} \end{aligned}$$

which demonstrates that the formula for the solution of (3.13)

is analogous to the formula (2.15) for the solution of (2.10).

Since $-\pi \cot \gamma\pi = \lambda$, the result (3.15) can also be written as

$$\begin{aligned} (3.16) \quad & (\tau-\alpha)^{1-\gamma} \phi(\tau) = - \frac{1}{\lambda^2 + \pi^2} \frac{d}{d\tau} \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\alpha\omega}} \frac{(\xi-\alpha)^{1-\gamma} f(\xi) d\xi d\omega}{(\omega-\xi)^{1-\gamma}} \\ & + \frac{c}{(\beta-\tau)^\gamma \frac{\lambda^2 + \pi^2}{\pi^2}} . \end{aligned}$$

The derivative of the double integral in (3.16) can be replaced by

$$\begin{aligned}
& \frac{d}{d\tau} \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\alpha\omega}} \frac{(\xi-\alpha)^{1-\gamma} f(\xi) d\xi d\omega}{(\omega-\xi)^{1-\gamma}} \\
&= \frac{d}{d\tau} \left\{ \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\alpha\tau}} \frac{(\xi-\alpha)^{1-\gamma} f(\xi) d\xi d\omega}{(\omega-\xi)^{1-\gamma}} \right. \\
&\quad \left. + \int_{L_{\tau\beta}} \frac{1}{(\omega-\tau)^\gamma} \int_{L_{\tau\beta}} \frac{\mu(\omega-\xi) \cdot (\xi-\alpha)^{1-\gamma} f(\xi) d\xi d\omega}{(\omega-\xi)^{1-\gamma}} \right\} \\
&= \frac{d}{d\tau} \left\{ \int_{L_{\alpha\tau}} (\xi-\alpha)^{1-\gamma} f(\xi) \int_{L_{\tau\beta}} \frac{(\omega-\xi)^{\gamma-1} d\omega d\xi}{(\omega-\tau)^\gamma} \right. \\
&\quad \left. + \int_{L_{\tau\beta}} (\xi-\alpha)^{1-\gamma} f(\xi) \int_{L_{\tau\beta}} \frac{(\omega-\xi)^{\gamma-1}}{(\omega-\tau)^\gamma} \cdot \mu(\omega-\xi) d\omega d\xi \right\} \\
&= \frac{d}{d\tau} \left\{ \int_{L_{\alpha\tau}} (\xi-\alpha)^{1-\gamma} f(\xi) \int_{L_{\tau\beta}} \frac{(\omega-\xi)^{\gamma-1} d\omega d\xi}{(\omega-\tau)^\gamma} \right. \\
&\quad \left. + \int_{L_{\tau\beta}} (\xi-\alpha)^{1-\gamma} f(\xi) \int_{L_{\xi\beta}} \frac{(\omega-\xi)^{\gamma-1} d\omega d\xi}{(\omega-\tau)^\gamma} \right\} \\
&= \frac{d}{d\tau} \left\{ \int_{L_{\alpha\tau}} (\xi-\alpha)^{1-\gamma} f(\xi) \int_{\infty}^{\frac{\beta-\xi}{\beta-\tau}} \frac{\sigma^{\gamma-1} d\sigma d\xi}{1-\sigma} \right. \\
&\quad \left. + \int_{L_{\tau\beta}} (\xi-\alpha)^{1-\gamma} f(\xi) \int_0^{\frac{\beta-\xi}{\beta-\tau}} \frac{\sigma^{\gamma-1} d\sigma d\xi}{1-\sigma} \right\}
\end{aligned}$$

$$\begin{aligned}
&= +(\tau-\alpha)^{1-\gamma} f(\tau) \int_0^{\infty} \frac{\sigma^{\gamma-1} d\sigma}{\sigma-1} \\
&\quad + \int_{L_{\alpha\beta}} (\zeta-\alpha)^{1-\gamma} f(\zeta) \left(\frac{\beta-\zeta}{\beta-\tau}\right)^{\gamma-1} \frac{1}{1 - \left(\frac{\beta-\zeta}{\beta-\tau}\right)} \cdot \frac{(\beta-\zeta)}{(\beta-\tau)^2} \cdot d\zeta \\
&= \lambda(\tau-\alpha)^{1-\gamma} f(\tau) + \frac{1}{(\beta-\tau)^\gamma} \int_{L_{\alpha\beta}} \frac{(\zeta-\alpha)^{1-\gamma} (\beta-\zeta)^\gamma f(\zeta) d\zeta}{\zeta-\tau} .
\end{aligned}$$

Hence (3.16) can be expressed as

$$\begin{aligned}
(3.17) \quad \phi(\tau) = & - \frac{\lambda f(\tau)}{\lambda^2 + \pi^2} - \frac{1}{(\lambda^2 + \pi^2)(\tau-\alpha)^{1-\gamma}(\beta-\tau)^\gamma} \int_{L_{\alpha\beta}} \frac{(\zeta-\alpha)^{1-\gamma} (\beta-\zeta)^\gamma f(\zeta) d\zeta}{\zeta-\tau} \\
& + \frac{c}{(\tau-\alpha)^{1-\gamma}(\beta-\tau)^\gamma \sqrt{\lambda^2 + \pi^2}}
\end{aligned}$$

where

$$-\pi \cot \gamma\pi = \lambda .$$

This is the form which is usually presented for the solution of

$$(3.18) \quad \int_{L_{\alpha\beta}} \frac{\phi(z) dz}{z-\zeta} = \lambda \phi(\zeta) + f(\zeta) .$$

It is to be noticed that although (3.17) depends only on a single integral the integrand contains the factor $1/(\zeta-\tau)$ and hence the integral denotes the Cauchy principal value. This factor does not appear in (3.16). For some purposes the formula (3.16) may be more advantageous than (3.17).

It is interesting to note that the foregoing analysis subsumes the solution of

$$(3.19) \quad \oint_C \frac{\phi(z)dz}{z-\zeta} = \lambda\phi(\zeta) + f(\zeta)$$

where C is a simple smooth closed path and ζ is on C . If α is any point on C it can be taken as both the initial and end point of a path $L_{\alpha\beta}$ which coincides with C . Hence the solution of (3.19) can be found by setting $\beta = \alpha$ in (3.16) or (3.17). The latter form gives

$$(3.20) \quad \phi(\tau) = -\frac{\lambda f(\tau)}{\lambda^2 + \pi^2} - \frac{1}{(\lambda^2 + \pi^2)(\tau - \alpha)} \oint_C \frac{(\zeta - \alpha)f(\zeta)d\zeta}{\zeta - \tau} \\ + \frac{c}{(\tau - \alpha)^{1-\gamma}(\alpha - \tau)^\gamma \sqrt{\lambda^2 + \pi^2}}$$

or

$$(3.21) \quad \phi(\tau) = -\frac{f(\tau)}{\lambda^2 + \pi^2} - \frac{1}{(\lambda^2 + \pi^2)} \oint_C \frac{f(\zeta)d\zeta}{\zeta - \tau} \\ - \frac{1}{(\lambda^2 + \pi^2)(\tau - \alpha)} \left[\oint_C f(\zeta)d\zeta - \frac{\pi e^{-\pi i \gamma}}{\sin \gamma \pi} \oint_C \phi(\zeta)d\zeta \right].$$

A solution of (3.19) must satisfy

$$\oint_C \oint_C \frac{\phi(z)dzd\zeta}{z-\zeta} = \lambda \oint_C \phi(\zeta)d\zeta + \oint_C f(\zeta)d\zeta$$

or

$$-\pi i \oint_C \phi(z) dz = \lambda \oint_C \phi(\xi) d\xi + \oint_C f(\xi) d\xi .$$

That is,

$$\oint_C f(\xi) d\xi = -(\lambda + \pi i) \oint_C \phi(\xi) d\xi$$

and since $-\pi \cot \gamma\pi = \lambda$,

$$\begin{aligned} \oint_C f(\xi) &= \pi(\cot \gamma\pi - i) \oint_C \phi(\xi) d\xi \\ &= \frac{\pi e^{-i\gamma\pi}}{\sin \gamma\pi} \cdot \oint_C \phi(\xi) d\xi \end{aligned}$$

which shows that the quantity within the bracket of (3.21) is zero. Therefore the solution of

$$\oint_C \frac{\phi(z) dz}{z - \xi} = \lambda \phi(\xi) + f(\xi)$$

is

$$\phi(\tau) = -\frac{\lambda f(\tau)}{\lambda^2 + \pi^2} - \frac{1}{(\lambda^2 + \pi^2)} \oint_C \frac{f(\xi) d\xi}{\xi - \tau} .$$

References

- [1] Peters, A. S., A Note on the Integral Equation of the First Kind with a Cauchy Kernel, Comm. Pure Appl. Math., Vol. XVI, No. 1, 1963, pp. 57-61.
- [2] Muskhelishvili, N. I., Singular Integral Equations, P. Noordhoff N. V., Groningen, Holland, 1953.
- [3] Mikhlin, S. G., Integral Equations, Pergamon Press, 1957.
- [4] Carrier, G. F.; Krook, M.; Pearson, C. E., Functions of a Complex Variable, McGraw-Hill Book Co., 1966.
- [5] Levinson, N., Simplified Treatment of Integrals of Cauchy Type, the Hilbert Problem and Singular Integral Equations. Appendix: Poincaré-Bertrand Formula, SIAM Review, Vol. 7, No. 4, 1965, pp. 474-502.
- [6] Carleman, T., Sur la Résolution de Certaines Équations Intégrales, Arkiv för Matematik Astronomi och Fysik, Bd. 16, No. 26, 1922.

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
Courant Institute of Mathematical Sciences New York University		not classified	
		2b. GROUP	
		none	
3. REPORT TITLE			
Abel's Equation and the Cauchy Integral Equation of the Second Kind			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Technical Report October 1966			
5. AUTHOR(S) (Last name, first name, initial)			
Peters, Arthur S.			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
October 1966		24	6
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
Nonr-285(55)		IMM 352	
b. PROJECT NO.			
NR 062-160			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		none	
10. AVAILABILITY/LIMITATION NOTICES			
Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
none		U.S. Navy, Office of Naval Research 207 West 24th St., New York, N.Y.	
13. ABSTRACT			
<p>This report shows that a Cauchy singular integral equation of the second kind can be solved by reducing it to Abel's integral equation. The reduction yields a new formula for the solution of the Cauchy equation.</p>			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

APPROVED DISTRIBUTION LIST

Chief of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 438 (3)	Chief, Bureau of Aeronautics Department of the Navy Washington 25, D. C. Attn: Research Division (1)
Commanding Officer Office of Naval Research Branch Office 219 S. Dearborn Street Chicago, Illinois 60604 (1)	Chief, Bureau of Ordnance Department of the Navy Washington 25, D. C. Attn: Research and Development Division (1)
Commanding Officer Office of Naval Research Branch Office 207 West 24th St. New York 11, N.Y. (1)	Office of Ordnance Research Department of the Army Washington 25, D. C. (1)
Commanding Officer Office of Naval Research Branch Office 1030 East Green Street Pasadena 1, Calif. (1)	Headquarters Air Research and Development Command United States Air Force Andrews Air Force Base Washington 25, D. C. (1)
(1)	Director of Research National Advisory Committee for Aeronautics 1724 F Street, Northwest Washington 25, D. C. (1)
Commanding Officer Office of Naval Research Box 39, Fleet Post Office New York, New York 09510 (5)	Director Langley Aeronautical Laboratory National Advisory Committee for Aeronautics Langley Field, Virginia (1)
Director Naval Research Laboratory Washington 25, D. C. Attn: Code 2021 (6)	Director National Bureau of Standards Washington 25, D. C. Attn: Fluid Mechanics Section (1)
Defense Documentation Center Cameron Station Alexandria, Va. 22314 (20)	Professor R. Courant Courant Institute of Mathematical Sciences, N.Y.U. 251 Mercer St. New York 12, N.Y. (1)
Professor W.R. Sears Director Graduate School of Aeronautical Engineering Cornell University Ithaca, New York (1)	Professor G. Kuertl Department of Mechanical Engineering Case Institute of Technology Cleveland, Ohio (1)

DISTRIBUTION LIST (CONT.)

Chief, Bureau of Ships
Department of the Navy
Washington 25, D. C.
Attn: Research Division (1)
Code 420 Preliminary
Design (1)

Commander
Naval Ordnance Test Station
3202 E. Foothill Blvd.
Pasadena, Calif. (1)

Commanding Officer and Director
David Taylor Model Basin
Washington 7, D. C.
Attn: Hydromechanics Lab. (1)
Hydrodynamics Div. (1)
Library (1)
Ship Division (1)

California Institute of
Technology
Hydrodynamic Laboratory
Pasadena 4, California (1)

Professor A.T. Ippen
Hydrodynamics Laboratory
Massachusetts Institute
of Technology
Cambridge 39, Mass. (1)

Dr. Hunter Rouse, Director
Iowa Institute of Hydraulic
Research
State University of Iowa
Iowa City, Iowa (1)

Stevens Institute of Technology
Experimental Towing Tank
711 Hudson Street
Hoboken, New Jersey (1)

Dr. G. H. Hickox
Engineering Experiment Station
University of Tennessee
Knoxville, Tennessee (1)

Dr. L. G. Straub
St. Anthony Falls Hydraulic
Laboratory
University of Minnesota
Minneapolis 14, Minn. (1)

Chief of Naval Research
Department of the Navy
Washington 25, D. C.
Attn: Code 416 (1)
Code 460 (1)

Chief, Bureau of Yards and Docks
Department of the Navy
Washington 25, D. C.
Attn: Research Division (1)

Hydrographer
Department of the Navy
Washington 25, D. C. (1)

Director
Waterways Experiment Station
Box 631
Vicksburg, Mississippi (1)

Office of the Chief of Engineers
Department of the Army
Gravelly Point
Washington 25, D. C. (1)

Beach Erosion Board
U.S. Army Corps of Engineers
Washington 25, D. C. (1)

Commissioner
Bureau of Reclamation
Washington 25, D. C. (1)

Dr. G. H. Keulegan
National Hydraulic Laboratory
National Bureau of Standards
Washington 25, D. C. (1)

Brown University
Graduate Division of Applied
Mathematics
Providence 12, Rhode Island (1)

California Institute of
Technology
Hydrodynamics Laboratory
Pasadena 4, California
Attn: Professor M. S. Plesset (1)
Professor V.A. Vanoni (1)

DISTRIBUTION LIST (CONT.)

Professor M. L. Albertson
Department of Civil Engineering
Colorado A. + M. College
Fort Collins, Colorado (1)

Professor G. Birkhoff
Department of Mathematics
Harvard University
Cambridge 38, Mass. (1)

Massachusetts Institute of
Technology
Department of Naval Architecture
Cambridge 39, Mass. (1)

Dr. R. R. Revelle
Scripps Institute of Oceanography
La Jolla, California (1)

Stanford University
Applied Mathematics and
Statistics Laboratory
Stanford, California (1)

Professor H.A. Einstein
Department of Engineering
University of California
Berkeley 4, Calif. (1)

Dean K. L. Schoenherr
College of Engineering
University of Notre Dame
Notre Dame, Indiana (1)

Director
Woods Hole Oceanographic
Institute
Woods Hole, Mass. (1)

Professor J.W. Johnson
Fluid Mechanics Laboratory
University of California
Berkeley 4, Calif. (1)

c. 2.

c.2

c.2

DATE DUE

NOV 13 89			
EDIC			
PLAC 2'			
14N 7-107E			
TEN			
GAYLORD			PRINTED NO. 4